Field-tested laboratory-derived models to predict forest fire front spread rate

Autor(es): Rossa, Carlos G.; Fernandes, Paulo M.

Publicado por: Imprensa da Universidade de Coimbra

URL persistente: URI:<http://hdl.handle.net/10316.2/44674>

DOI: DOI:https://doi.org/10.14195/978-989-26-16-506_157

Accessed : 26-Apr-2023 11:32:51

A navegação consulta e descarregamento dos títulos inseridos nas Bibliotecas Digitais UC Digitalis, UC Pombalina e UC Impactum, pressupõem a aceitação plena e sem reservas dos Termos e Condições de Uso destas Bibliotecas Digitais, disponíveis em <https://digitalis.uc.pt/pt-pt/termos>.

Conforme exposto nos referidos Termos e Condições de Uso, o descarregamento de títulos de acesso restrito requer uma licença válida de autorização devendo o utilizador aceder ao(s) documento(s) a partir de um endereço de IP da instituição detentora da supramencionada licença.

Ao utilizador é apenas permitido o descarregamento para uso pessoal, pelo que o emprego do(s) título(s) descarregado(s) para outro fim, designadamente comercial, carece de autorização do respetivo autor ou editor da obra.

Na medida em que todas as obras da UC Digitalis se encontram protegidas pelo Código do Direito de Autor e Direitos Conexos e demais legislação aplicável, toda a cópia, parcial ou total, deste documento, nos casos em que é legalmente admitida, deverá conter ou fazer-se acompanhar por este aviso.

ADVANCES IN FOREST FIRE RESEARCH 2018

EDITED BY

DOMINGOS XAVIER VIEGAS

ADAI/CEIF, UNIVERSITY OF COIMBRA, PORTUGAL

Short contribution – Fire Management

Field-tested laboratory-derived models to predict forest fire front spread rate

Carlos G. Rossa*; Paulo M. Fernandes

*Centre for the Research and Technology of Agro-environmental and Biological Sciences (CITAB),
University of Trás-os-Montes e Alto Douro (UTAD). Quinta de Prados, Apartado 1013, 5001-801 Vila
Real, Portugal. {carlos.g.rossa@gmail.com*, pfern@utad.pt}*

Keywords: fire behaviour, live fuels, empirical models, outdoors fires, fire management.

1. Introduction

Forest fire-spread rate models provide crucial support to decision support in fire management, such as assisting wildfire suppression operations. Most modelling studies aim at predicting fire spread rate (R) for wind- or slope-driven fires (Sullivan et al. 2014). Even so, the ability to estimate basic fire spread rate (R_0), i.e., on level ground in the absence of wind, is relevant because it provides a measure of fuel bed flammability and some modelling approaches estimate R by multiplying R_0 by a wind or slope factor (e.g., Rothermel 1972). Conversely, wind-driven fire spread rate (R_U) has been the motivation of many modelling efforts. Slope has also a strong effect on fire spread. However, when the distance travelled is sizably larger than distances between landscape undulations, R is satisfactorily equated to that of a fire spreading across a corresponding expanse of flat ground (Sullivan et al. 2014). As a result, the distance travelled by large fires is mainly a function of wind speed. Yet, a field-tested empirical R_U model for fire spread in generic vegetation is currently lacking.

In this work, we present results of laboratory experiments to analyse fire spread under R_0 and R_U conditions. The data are used in the development of models for fire propagation in generic fuel beds. Fuel beds were built from natural vegetation, mimicking the real-world structure of litter and shrubs. Empirical models were derived using regression analysis.

2. Data and methods

We conducted laboratory fire spread experiments in litter and shrub-like fuel beds. Some of the tests to measure R_0 ($n = 220$) were immediately followed by a R_U test in an identical fuel bed ($n = 108$), to isolate and assess the effect of wind. Shrub-like fuel beds were composed of vertically-placed tree or shrub twigs, sometimes under-layered by litter (Figure 1). We used various plant species (*Pinus pinaster* Ait., *Eucalyptus globulus* Labill., *Eucalyptus obliqua* L'Her., *Acacia mangium* Willd., *Quercus robur* L., *Pinus resinosa* Sol. ex Ait.) and fuel beds were often composed of mixed dead and quasi-live vegetation. Quasi-live fuels were collected live, with fuel moisture content (M) decreasing as a function of storage time. R and M were measured in all tests and flame length and angle were also assessed in some of the trials. All fuel beds were at least 1-m long and 1 to 1.2-m wide.

We used regression analysis to relate R_0 with the pertinent variables. The effect of wind was assessed the same way and the wind effect and R_0 were multiplied to predict R_U . The laboratory-derived relationships were then tested against independent laboratory and field fires. Independent datasets were retrieved from the literature and included fire spread in no-wind and no-slope conditions, as well as wind-driven fires.

Figure 1 - Wind-driven fire spread test in a fuel bed of Eucalyptus globulus leaves litter over layered by quasi-live E. globulus twigs.

3. Initial results and conclusion

Our results show that both R_0 and R_U models can be predicted by a single formulation and a common set of variables, independently of vegetation type. Although the models were fitted to laboratory experiments data, they were tested against independent laboratory and field fires data, and hence justify further testing in other real-world fire spread situations for a wider range of fuel complexes.

4. Acknowledgments

This work was supported by Fundação para a Ciência e a Tecnologia (FCT) under post-doctoral grant SFRH/BPD/84770/2012 (financing programs POPH and FSE) to the first author. It was also supported by FCT and FEDER – Fundo Europeu de Desenvolvimento Regional (COMPETE 2020, POCI – Programa Operacional Competitividade e Internacionalização), in the framework of projects BONFIRE (POCI-01-0145-FEDER-016727, PTDC/AAGMAA/2656/2014) and CITAB (UID/AGR/04033/2013, POCI-01-0145-FEDER-006958). We are grateful to Délio Sousa, Paulo Ventura, and André Rosas (University of Trás-os-Montes e Alto Douro) for assisting in some of the laboratory experimental work. The help of Fernando Frazão, David Davim, José Silva, Nuno Marcos, Rafael Figueiredo, and Raquel Roliz in fuel collection is greatly appreciated.

5. References

- Rothermel RC (1972) A mathematical model for predicting fire spread in wildland fuels. USDA Forest Service, Intermountain Forest and Range Experiment Station, Research Paper INT-115. (Ogden, UT)
- Sullivan AL, Sharples JJ, Matthews S, Plucinski M (2014) Estimating downslope fire spread for landscape fire spread prediction. *Environmental Modelling & Software* 62, 153–163. doi:10.1016/j.envsoft.2014.08.024