The spawning migration of the sea lamprey (*Petromyzon morinus* L.) in the river Mondego

Autor(es): Almeida, Pedro Raposo; Silva, Pedro Teixeira da; Quintela, Bernardo Ruivo

Publicado por: Imprensa da Universidade de Coimbra

URL persistente: URI:<http://hdl.handle.net/10316.2/32707>

DOI: DOI:http://dx.doi.org/10.14195/978-989-26-0336-0_25

Accessed : 26-Apr-2023 05:32:59

A navegação consulta e descarregamento dos títulos inseridos nas Bibliotecas Digitais UC Digitalis, UC Pombalina e UC Impactum, pressupõem a aceitação plena e sem reservas dos Termos e Condições de Uso destas Bibliotecas Digitais, disponíveis em <https://digitalis.uc.pt/pt-pt/termos>.

Conforme exposto nos referidos Termos e Condições de Uso, o descarregamento de títulos de acesso restrito requer uma licença válida de autorização devendo o utilizador aceder ao(s) documento(s) a partir de um endereço de IP da instituição detentora da supramencionada licença.

Ao utilizador é apenas permitido o descarregamento para uso pessoal, pelo que o emprego do(s) título(s) descarregado(s) para outro fim, designadamente comercial, carece de autorização do respetivo autor ou editor da obra.

Na medida em que todas as obras da UC Digitalis se encontram protegidas pelo Código do Direito de Autor e Direitos Conexos e demais legislação aplicável, toda a cópia, parcial ou total, deste documento, nos casos em que é legalmente admitida, deverá conter ou fazer-se acompanhar por este aviso.

MIGUEL ÂNGELO PARDAL
JOÃO CARLOS MARQUES
MANUEL AUGUSTO GRAÇA
Scientific Editors

Aquatic Ecology of the Mondego River Basin Global Importance of Local Experience

Coimbra • Imprensa da Universidade

MIGUEL ÂNGELO PARDAL
JOÃO CARLOS MARQUES
MANUEL AUGUSTO GRAÇA
Scientific Editors

Aquatic Ecology of the Mondego River Basin Global Importance of Local Experience

Coimbra • Imprensa da Universidade

COORDENAÇÃO EDITORIAL
Imprensa da Universidade de Coimbra

CONCEPÇÃO GRÁFICA
António Barros

INFOGRAFIA
António Resende
Estímulus [design] • Coimbra

EXECUÇÃO GRÁFICA
GRAFIASA

ILUSTRAÇÃO DA CAPA
P. P. Cunha e J. Dinis

ISBN
972-8704-04-6

DEPÓSITO LEGAL
175038/02

© JANEIRO 2002, IMPRENSA DA UNIVERSIDADE DE COIMBRA

OBRA PUBLICADA COM O PATROCÍNIO DE:
IMAR – INSTITUTO DO MAR
IPIMAR – INSTITUTO DE INVESTIGAÇÃO DAS PESCAS E DO MAR

PEDRO RAPOSO ALMEIDA ^{1,2}
HUGO TEIXEIRA DA SILVA ¹
BERNARDO RUIVO QUINTELLA ¹

THE SPAWNING MIGRATION OF THE SEA LAMPREY (*Petromyzon marinus* L.) IN THE RIVER MONDEGO

Abstract

Presently, sea lampreys (*Petromyzon marinus* L.) are confined to the last 35 km of the River Mondego, downstream from the Açude-Ponte dam. Their spawning migration occurs between December and May-June, with a peak in February-March. During that period, this cyclostome presents a nocturnal behaviour, performing their upstream migration during the night, dawn and early morning periods.

Although the physical obstacles present in the freshwater zone of River Mondego retard the upstream movement, the submerged stone weirs create riffles, used by the sea lampreys as resting areas during the day.

The concentration of nests is higher in the first 2 km downstream from the Açude-Ponte dam. After hatching, the ammocoetes spend four years in freshwater habitats, the higher densities being located near the nest area.

Apart from habitat destruction, resulting from dam construction and channelisation, intense fishing and poaching are the major threats to the survival of the sea lampreys in the River Mondego.

Introduction

The River Mondego watershed is one of the most important Portuguese river basins for the diadromous fish. Four species are commercially exploited in this region, the sea lamprey (*Petromyzon marinus* L.), the allis shad (*Alosa alosa* L.), the twaite shad (*A. fallax* Lacépède) and the eel (*Anguilla anguilla* L.).

Assis et al. (1992) point out that the economic value of the diadromous species is directly related with the predictability of the migratory season and movement route, as well as with the occurrence of large concentrations of animals in a short time-space scale, which makes them an easy target for fishermen and natural predators.

⁽¹⁾ Instituto de Oceanografia, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

⁽²⁾ Laboratório de Biologia da Conservação, Departamento de Biologia, Universidade de Évora, 7002-554 Évora, Portugal

The sea lamprey is a migratory species whose anadromous form is exploited as a commercial fishery in most of the major rivers in Portugal, particularly in the Central and Northern regions of the country. Sea lampreys are mostly captured when entering the estuaries to initiate their upstream reproductive migration, corresponding to the terminal phase of their life cycle (Beamish 1980, Ducasse and Leprince 1980, Guimarães 1987). As the species is considered a delicacy, its commercial value is rather high, attaining market values of 9500 PTE ($\cong$ € 47) per unit.

P. marinus is considered a "Vulnerable" species in the Portuguese Red Book of Endangered Species (Vários 1991), and the literature is unanimous in stating that the sea lamprey's populations are decreasing in all major Portuguese rivers (Guimarães 1987, Alexandrino 1990, Almaca 1990, Assis et al. 1992, Assis 1994, Ferreira and Oliveira 1996).

The human activities that promote significant changes in the physical and chemical characteristics of running water ecosystems can have major impacts in the sea lamprey population dynamics. Several authors have stressed the direct and indirect impacts on the conservation of diadromous species in general, and on the sea-lamprey in particular, resulting from the construction of dams, land recovery (drainage and flood protection), pollution (agricultural, forestry, industrial and urban use), dredging and gravel extraction, overfishing and poaching (Ducasse and Leprince 1980, Assis et al. 1992, Witkowski 1992, Young et al. 1990).

This paper describes the spawning migration of the sea lamprey in River Mondego, based on information gathered from previous works, and from the data obtained by the authors in a recent research project, involving the study of the migratory behaviour observed by ultrasonic tracking in estuarine and freshwater environments. The major threats to the survival of this species in the River Mondego watershed are identified and discussed.

The spawning migration and freshwater phase of the sea lamprey life cycle

Before dam construction, anadromous fish used to migrate into the upper reaches of River Mondego, entering some of the tributaries, namely the rivers Ceira and Alva. Since the construction of the Açude-Ponte dam, at Coimbra, in the early 80's, the migratory species are unable to pass this obstacle, and are forced to complete their life cycle in the last 35 km of the river (Fig. 1). This lower portion of the river runs in open valleys, in a plain area, has two main tributaries, the rivers Arunca and Pranto, and can be divided into two ecologically different stretches: an estuarine, and a freshwater environment. The separation between these two zones results from the presence of a 3 m height blockstone weir, the Formoselha weir, located 11 km downstream from the Açude-Ponte dam. The tidal effect is no longer visible upstream from this weir, and therefore the estuary is limited to the final 24 km of the river. The 11 km freshwater stretch between the Formoselha weir and the Açude-Ponte dam has 11 riffle areas, resulting from partially submerged stone weirs (transversal sediment retention platforms) installed across the river to avoid downstream sediment transportation.

Figure 1. Schematic representation of the River Mondego downstream from the Açude-Ponte dam (adapted from Almeida et al. 2000)

The professional fishermen's activity is restricted to two river stretches (Fig. 1) and is only allowed between the 15 January and the 15 April (Anonymous 1972, 1988).

Although the spawning migration of *P. marinus* in the River Mondego starts in late December (Guimarães 1987, Almeida et al. 2000), the period of time spent by these animals in the estuary or in the coastal vicinity, before they engage in their upstream migration, is still unknown. The migration continues until May-June, with a peak in February-March (Guimarães 1987, Alexandrino 1990, Machado-Cruz et al. 1990, Almeida et al. 2000). According to the information obtained from local fishermen, some spawning activity can still be observed in early August.

During their spawning migration, sea lampreys entering River Mondego present a daily migratory behaviour characterised by an alternation between resting periods and continuous upstream movements Almeida et al. (2000). According to these authors, the dusk hours are also associated with an increase in the activity of sea lampreys, but the upstream migration occurs during the night, dawn and early morning periods (i.e. between one hour after sunset to 9 a.m.). This nocturnal behaviour has already been observed by Hardisty and Potter (1971), Hardisty (1979, 1986) and Stier and Kynard (1986).

In the studies performed in the River Mondego, Almeida et al. (2000) found that both in the estuarine and in the freshwater environments, the movements of the sea lampreys present identical speeds. The apparently better performances showed by the animals in the estuary probably resulted from the existence of longer river stretches free from physical obstacles, enabling them to present a cruising ground speed of 16.5 body-lengths min⁻¹. This migratory behaviour can be kept for, approximately, nine hours of continuous movement (Almeida et al. 2000).

In freshwater, longer periods of continuous movement resulted in a reduction in the movement speed. The presence of numerous obstacles, namely, sand banks, shallow water sections, riffles and aquatic vegetation, significantly increased the difficulty to reach the upstream stretches of the river (Almeida et al. 2000)

The submerged stone weirs, present in the freshwater section between the Açude-Ponte dam and the Formoselha weir, are responsible for a considerable delay in the upriver progression of the sea lampreys and consequently for an extra energy

consumption. On the other hand, these weirs provide the necessary disruption in the habitat continuity, with the formation of riffles, used by these animals as preferential resting areas during the day (Almeida et al. 2000).

The spawning migration ends with the arrival at the spawning site, nest building, and finally, matting and fertilisation. Nest building is mainly performed by males, usually the first to arrive at the spawning sites (Hardisty and Potter 1971, Hardisty 1979). However, Hardisty and Potter (1971) refer that, late in the spawning season, females may assist males in the construction of the nests.

The choice of spawning sites may be influenced by some abiotic factors, specially the existence of an unidirectional water flow, and the presence of a substrate with adequate characteristics to the nest construction (coarse gravel, pebbles and sand) (Ducasse and Leprince 1980, Manion and Hansen 1980, Hardisty 1986). In the River Mondego, besides these factors, the existence of a barrier that cannot be transposed, the Açude-Ponte dam, plays an important role in determining the concentration of nests in the 2 km downstream from the Açude-Ponte dam. These nests are usually located in relatively deep water (50 cm), in the river stretches between the submerged stone weirs.

Males are the first to arrive at the spawning grounds, establishing and protecting a nesting territory (Manion and Hansen 1980). In the River Mondego, poachers take advantage of this behaviour to capture the animals while they are engaged in mating activities (unpublished data).

Like other lampreys, *P. marinus* spends most of its life in freshwater environment (migratory phase and larval stage). Hardisty (1979) believes that the marine phase of the lampreys' life cycle representing only 1/3 to 1/4 of the animal's life, is probably a secondary introduction in the life cycle of a group which primarily lived in freshwater environments.

Figure 2. Sea lamprey (*P. marinus* L.) life cycle (adapted from Hardisty and Potter 1971).

The duration of this larval stage depends on the abiotic conditions of the river, and varies between 4 to 7 years (Hardisty and Potter 1971, Sousa 1988, Afonso 1989) (Fig. 2). In the case of the River Mondego, the ammocoetes spend, at least four years in freshwater habitats, before they undergo a process of metamorphosis that will lead them through a macroptalmic stage to a juvenile form, identical to the adult (Quintella 2000). As a result, the success of a lamprey population is highly dependent on the quality of the freshwater environment, which should provide the appropriate conditions to ensure the spawning success, the development of the eggs and the survival of the ammocoetes.

The authorities responsible for the management of the River Mondego watershed should keep in mind the peculiarities of the sea lampreys' migratory behaviour. From December to June, and at least during the period between dusk and early morning (i.e. 1 h before sunset to 3 h after sunrise), the freshwater flow should allow the sea lampreys to reach the upstream spawning grounds with less effort. Furthermore, it is important to maintain the riffle areas. However, some of the physical obstacles (e.g. extensive shallow areas) present in the freshwater zone, upstream from the Formoselha weir, should be eliminated.

Downstream the Açude-Ponte dam the major impacts of man's activities on the aquatic environment result from agriculture-generated wastes, dam construction and channelisation. Among these three factors, dam construction induces the worst consequences to the sea lamprey population. The blocking of longitudinal connectivity drastically reduced the river basin area accessible to the migratory species. Besides, the disturbance of flow regimes downstream from the dams can attenuate the abiotic cues that are usually associated with the entering of the adult sea lampreys in a specific river (Hardisty 1979).

Although eutrophication caused by agriculture-generated wastes and domestic sewage could promote disturbances in the riverine ecosystem, the resulting increase in the phytoplankton production and in the accumulation of particulate organic matter in the sediment, might have been responsible for the good condition factor and fast growth rate of the ammocoete populations (Quintella 2000).

Finally, but by no means less important, the fishing effort from professional fishermen and poachers should be reduced, at least while there is no scientific evidence that the sea lamprey population is recovering.

Acknowledgements

Thanks are due to our colleague Dr. C.A. Assis for the useful comments on early drafts of the typescript.

References

- Afonso, A.M.S. 1989. A lampreia-mancha (*Petromyzon marinus*): aspectos morfológicos da metamorfose e sua relação com a biologia da espécie. Provas de Aptidão Pedagógica e Capacidade Científica, University of Oporto, Oporto.

- Alexandrino, P.J.B. 1990. Dispositivos de transposição de barragens para peixes migradores, em deslocação para montante. Provas de Aptidão Pedagógica e Capacidade Científica, University of Oporto, Oporto.
- Almaça, C. 1990. A lampreia e o estuário na bacia do Douro. Observatório, 1: 377-382.
- Almeida, P.R., Silva, H.T. and Quintella, B.R. (2000). The migratory behaviour of the sea lamprey *Petromyzon mannus* L., observed by acoustic telemetry in River Mondego (Portugal). In: A. Moore and Russel I. (eds.) Advances in fish telemetry. CEFAS, Lowestoft, Suffolk.
- Anonymous 1972. Decreto-Lei 265/72. Ministério da Marinha. Direcção-Geral dos Serviços de Fomento Marítimo. Diário do Governo, Série I. 177, 31.
- Anonymous 1988. Portaria 47/88. Ministério da Agricultura, Pescas e Alimentação. Secretaria de Estado da Agricultura. Diário da República, Série I. 19, 258.
- Assis, C.A. 1994. Migrações e peixes migradores. Uma perspectiva conservacionista. O Biólogo 29:13-16.
- Assis, C.A., Costa, J.L., Costa, M. J., Moreira, F., Almeida, P. and Gonçalves, J.J. 1992. Ameaças à sobrevivência dos peixes migradores do Tejo. Sugestões para a sua conservação. Publicações Avulsas do INIP 17: 429-441.
- Beamish, F.W.H. 1980. Biology of the North American anadromous sea lamprey, *Petromyzon mannus*. Canadian Journal of Fisheries and Aquatic Sciences 37: 1924-1943.
- Ducasse, J. and Leprieux, Y. 1980. Étude préliminaire de la biologie des lamproies dans les bassins de la Garonne et de la Dordogne. Memoire. École Nationale des Ingenieurs des Travaux des Eaux and Forêts. 136p.
- Ferreira, M.T. and Oliveira, J.M. 1996. Gestão da lampreia-mancha *Petromyzon marinus* no rio Tejo. Anuário do Instituto Superior de Agronomia 45: 401-439.
- Guimarães, M.T. 1987. Medidas para a protecção e fomento da lampreia-do-mar (*Petromyzon marinus* L.) no rio Mondego. Actas do Colóquio Luso-Espanhol sobre Ecologia das Bacias Hidrográficas e Recursos Zoológicos, 195-203.
- Hardisty, M.W. 1979. Biology of Cyclostomes. Chapman and Hall, London. 428p.
- Hardisty, M.W. 1986. *Petromyzon marinus* Linnaeus, 1758. In The freshwater fishes of Europe. Petromyzontiformes. Edited by J. Holcik. Aula-Verlag Wiesbaden, Darmstadt. pp. 94-116.
- Hardisty, M.W. and Potter, I.C. 1971. The behaviour, ecology and growth of larval lampreys. In The Biology of Lampreys. Edited by M.W. Hardisty and I.C. Potter). Academic Press, London. pp. 85-125.
- Machado-Cruz, J.M., Valente, A.C.N. and Alexandrino, P.J.B. 1990. Contribuição para a caracterização ecológica e económica da pesca de migradores a jusante da barragem de Belver, Rio Tejo. Actas 1º Congresso do Tejo. Que Tejo, que futuro?: 189-200.
- Manion, P.J. and L.H. Hansen, 1980. Spawning behaviour and fecundity of lampreys from the upper three Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 37: 1635-1640.
- Sousa, J.A. 1988. Dados preliminares sobre a biologia da lampreia, *Petromyzon mannus*, no rio Lima. Actas do Colóquio Luso-Espanhol sobre Ecologia das Bacias Hidrográficas e Recursos Zoológicos: 405-414.
- Stier, K. and Kynard, B. 1986. Movement of sea-run sea lampreys, *Petromyzon marinus*, during the spawning migration in the Connecticut river. Fishery Bulletin 84: 749-753.
- Quintella, B.R. 2000. Ecologia da fase larvar de (*Petromyzon marinus* L.) no Baixo Mondego. Relatório de Estágio. University of Lisbon, Lisbon. 48p.
- Vários 1991. Livro Vermelho dos Vertebrados de Portugal. Vol. II - Peixes Dulciaquícolas e Migradores. SNPRCN, Lisboa. 55p.
- Witkowski, A., 1992. Threats and protection of freshwater fishes in Poland. Netherlands Journal of Zoology 42: 243-259.
- Young, R.J., Houston, K.A., Weise, J.G. and Kelso, J.R.M. 1990. The effect of environmental variables on the population dynamics of sea lamprey, *Petromyzon mannus*. Canadian Technical Reports on Fisheries and Aquatic Sciences, No. 1736.

Série

Investigação

•

Coimbra
Imprensa da Universidade

2002