

ANNALS OF RESEARCH IN SPORT AND PHYSICAL ACTIVITY

The effect of warm-up in short distance swimming performance

Autor(es): Neiva, Henrique P.; Marques, Mário C.; Bacelar, Lara; Moínhos, Nuno; Morouço, Pedro G.; Marinho, Daniel A.

URL persistente: URI:<http://hdl.handle.net/10316.2/5686>

DOI: DOI:http://dx.doi.org/10.14195/2182-7087_3_4

Accessed : 2-Jul-2022 09:33:31

A navegação consulta e descarregamento dos títulos inseridos nas Bibliotecas Digitais UC Digitalis, UC Pombalina e UC Impactum, pressupõem a aceitação plena e sem reservas dos Termos e Condições de Uso destas Bibliotecas Digitais, disponíveis em <https://digitalis.uc.pt/pt-pt/termos>.

Conforme exposto nos referidos Termos e Condições de Uso, o descarregamento de títulos de acesso restrito requer uma licença válida de autorização devendo o utilizador aceder ao(s) documento(s) a partir de um endereço de IP da instituição detentora da supramencionada licença.

Ao utilizador é apenas permitido o descarregamento para uso pessoal, pelo que o emprego do(s) título(s) descarregado(s) para outro fim, designadamente comercial, carece de autorização do respetivo autor ou editor da obra.

Na medida em que todas as obras da UC Digitalis se encontram protegidas pelo Código do Direito de Autor e Direitos Conexos e demais legislação aplicável, toda a cópia, parcial ou total, deste documento, nos casos em que é legalmente admitida, deverá conter ou fazer-se acompanhar por este aviso.





**ANNALS OF
RESEARCH IN SPORT
AND
PHYSICAL ACTIVITY**

FACULDADE DE
CIÊNCIAS DO
DESPORTO E
EDUCAÇÃO FÍSICA
DA UNIVERSIDADE
DE COIMBRA

IMPRENSA
DA UNIVERISDADE
DE COIMBRA

2 • SCIENCE AND SWIMMING

THE EFFECT OF WARM UP IN SHORT DISTANCE SWIMMING PERFORMANCE

Henrique P. Neiva^{1,2}, Mário C. Marques^{1,2}, Lara Bacelar^{1,2},
Nuno Moinhos^{1,2}, Pedro G. Morouço^{2,3}, Daniel A. Marinho^{1,2}

RUNNING HEAD: Warm-up effect in short distance swimming performance

ABSTRACT

Warming up before physical activity is usual and became assumed as essential in competition and training events. It is expected an optimization in performance but the literature is still ambiguous on this subject. The aim of this study was to assess the effect of the regular warm-up in 50 m swimming performance, in female swimmers. Seven national-level swimmers (mean \pm SD; age 15.3 ± 1.1 years-old, height: 1.61 ± 8.1 m, body mass: 56.5 ± 7.0 kg) volunteered for this study. Each swimmer performed 50 m freestyle at the maximum velocity, after previous warm-up and without performing the same, with 24 h between conditions. Times were registered and capillary blood lactate concentration was assessed after the swimming trial at the 1st and 3rd min of recovery. Additionally, the Borg ratings of perceived exertion scale were used and biomechanical parameters such as stroke frequency, stroke length and stroke index were assessed. The 50 m swimming times were not different with and without warm-up (33.05 ± 2.34 s and 32.71 ± 2.07 s, respectively, $p = .40$). No differences were found in lactate values (8.63 ± 1.49 mmol·l⁻¹ and 7.93 ± 1.92 mmol·l⁻¹, respectively; $p = .71$), ratings of perceived exertion (15.86 ± 1.07 and 15.14 ± 1.22 , respectively; $p = .24$), stroke frequency (0.81 ± 0.08 Hz and 0.81 ± 0.04 Hz, respectively; $p = .79$), stroke length (1.87 ± 0.14 m and 1.89 ± 0.12 m, respectively; $p = .74$) and stroke index (2.85 ± 0.31 m² c⁻¹ s⁻¹ and 2.91 ± 0.34 m² c⁻¹ s⁻¹, respectively; $p = .40$). These results suggested that regular warm-up used by the swimmers does not influence the 50 m freestyle performance, in female swimmers.

KEYWORDS: evaluation; female; freestyle; lactate; biomechanics.

INTRODUCTION

Warm-up is commonly known as a preparatory practice that aimed to improve the subsequent performance (Hedrick, 1992). It is intended to result in an improvement of muscle dynamics, reduce the risk of injury and prepare the athlete for the demands of the main activity

(Woods, Bishop & Jones, 2007). This positive effect on athletic performance seems to be a widespread belief of coaches and their athletes, though the scarcity of conclusive scientific evidence (Atkinson, Todd, Reilly & Waterhouse, 2005; Burnley, & Jones, 2002). When the muscle is exposed to physical activity, there is an increase in their temperature, proportionally to the work done (Saltin, Gage, & Stolwijk, 1968). The mechanisms related to the raise of core and muscle temperature appear to be of great importance for the proposed effects of warming-up before physical activity (Bishop, 2003a). Muscular performance is affected by decreasing the viscous resistance of the muscles and joints (Wright, 1973) and decreasing stiffness of the fibers during muscle contraction (Buchthal, Kaiser, Knappeis & 1944). Complementary, it is suggested to increase nerve conduction rate and speeding of metabolic reactions, such as the muscle glycogenolysis, glycolysis and high-energy phosphate degradation (Febbraio et al., 1996). Moreover, the hyperthermia resultant of the activity might contribute to increase the blood flow (Pearson et al., 2011) and the oxygen delivery to the muscles, via a rightward shift in the oxyhaemoglobin dissociation curve and the vasodilatation of muscle blood vessels (McCutcheon et al., 1999). Warm-up exercises also allowed the athletes to begin subsequent tasks with an elevated baseline of oxygen consumption, saving more anaerobic capacity for later in the task (Febbraio et al., 1996). Although all these effects proposing improvements in performance after the warm-up routines, the specialized literature is not clear in this matter. There were several studies demonstrating improvements in performance after warming-up (i.e., Atkinson et al., 2005; Burnley et al., 2002), and there were others reporting no changes or even detrimental changes in performance (i.e., Mitchell and Huston, 1993; Bishop, Bonetti, & Dawson, 2001).

Specifically in swimming, the few existing studies are unclear regarding to changes in performance. Moreover, some of them were restricted to physiological assessment, omitting the performance values and limiting the conclusions (Houmar et al., 1991; Robergs et al 1990). DeVries (1959) and Thompson (1958) suggested improvements in swimming velocity in distances until 91 m. These results were confirmed by Romney and Nethery (1993), who were able to verify significant improvements in 100 yards performance after 15 min of warm-up when compared with the absence of any activity prior to the trial. More recently, the maximal and mean values of propulsive force during 30 s of tethered swimming appeared enhanced after regular warm-up (Neiva, Morouço, Silva, Marques, & Marinho, 2011). If, on one hand these results reinforce the general idea that the warm-up is beneficial to the swimmer performance, there are several studies contributing to the controversy on this subject.

No effects in performance were suggested by Mitchell and Huston (1993), and Bobo (1999) in 100 m swimming trials. Regarding to short distance swimming, Neiva, Morouço, Pereira, and Marinho (2012) revealed that regular warm-up did not improve 50 m swimming performance. Blood lactate concentration ([La-]) and perceived exertion were also similar between no warm up and warm up conditions. These results are the opposite of those verified in the study of Balilionis et al. (2012). Better swimming times in 50 yards (0.2 s) after

regular warm-up procedures were revealed. Therefore, the contradictory results emerges the need to more investigation to further determine the influence of warm-up procedures, their optimal structure and specificity related to each sport. Hence, the aim of the current study was to verify the effect of warm-up procedures in short distance swimming performance (50 m freestyle), in female swimmers.

This way, it is intended to better understand and better know the effects of warming up in swimming.

METHODS

Subjects

Seven female swimmers (mean \pm SD; age: 15.3 ± 1.1 year-old, height: 1.61 ± 8.1 m, body mass: 56.5 ± 7.0 kg, fat mass: 14.3 ± 3.3 kg) participated in this study. Body mass and fat mass were assessed through a bioelectric impedance analysis method (Tanita BC 420S MA, Japan). All of them participate in the National Championships and their training experience was of 5 years at least. Volunteer's parents and coaches were informed about research propose and signed an informed consent to participate. All the procedures were approved by the institutional review board.

Procedures

All testing procedures were performed in a 50 m indoor swimming pool at a water temperature of 27.5 °C. The experiments took place one week after the main competition (National Championships) of the season second macrocycle. Each swimmer performed 50 m freestyle at the maximum velocity, in two different days. The use or no use of warm up procedures before the maximal test was the variable that distinguished two implemented protocols, performed with 24 hours of difference between them. In the warm-up condition, swimmers performed their usual warm-up before a competitive swimming event (1000 m of total volume). In-water starts were used. Times were registered by 2 experienced coaches using stopwatches (Seiko, Japan) and the average of these times was considered for further analysis. Capillary blood samples were collected from the fingertip before and after each swimming maximal test (at the 1st and 3rd min of recovery) to assess the higher value of [La-] (Accutrend Lactate®Roche, Germany). The Borg (1998) ratings of perceived exertion (RPE) scale of 15 points (6 – 20) was used to quantify exercise level of exertion after each test.

For biomechanical assessment both stroke frequency (SF) and stroke length (SL) were measured. SF was obtained with a chrono-frequency meter (Golfinho Sports MC 815, Aveiro, Portugal) from three consecutive stroke cycles, in the middle of the trial test. Then, SF values were converted to International System Units (Hz). SL was estimated as being (Craig, Skehan, Pawelczyk, & Boomer, 1985):

$$SL = \frac{V}{SF}$$

(1)

Where SL is the stroke length, V is the average velocity of the swimmer during the 50 m, and SF is the stroke frequency (Hz). The SI, considered as one of the swimming stroke efficiency indexes, was adapted and computed as (Costill, Kovaleski, Porter, Fielding, & King, 1985):

$$SI = V \times SL$$

(2)

Where SI is the stroke index ($m^2 c^{-1} s^{-1}$), V is the swimmer velocity ($m s^{-1}$) and the SL is the stroke length ($m c^{-1}$) of the swimmer.

Statistical analysis

88 • Standard statistical methods were used for calculation of means and standard deviations. The sample normality was determined by Shapiro-Wilk test. Since the very low value of the N (<30) and the rejection of the null hypothesis (H0), non-parametric procedures were adopted. In order to compare the data obtained with and without warm-up, nonparametric Wilcoxon signed rank test was used. Differences were considered significant for $p \leq 0.05$.

RESULTS

Table 1 presents the mean \pm SD values for the 50 m maximal swimming test variables assessed, namely the 50 m times and its partials of 25 m, blood lactate concentration, ratings of perceived exertion, stroke frequency, stroke length and stroke index. No significant differences were evident for the data obtained when comparing the warm-up condition with the no warm-up condition.


TABLE 1 Mean \pm standard deviation (mean \pm SD) values of 50 m swim time and each partial of 25 m (s), blood lactate concentration (mmol·l⁻¹), RPE, stroke frequency (Hz), stroke length (m.c-1) and stroke index (m² c⁻¹ s⁻¹) in the 50 m maximal test with no warm-up and with previous regular warm-up. *P* – values are presented.

	No WU Mean \pm SD	Regular WU Mean \pm SD	Sig (<i>p</i>)
1 st 25 m time (s)	15.01 \pm 0.90	15.24 \pm 1.06	0.31
2 nd 25 m time (s)	17.70 \pm 1.18	17.80 \pm 1.33	0.73
50 m time (s)	32.71 \pm 2.07	33.05 \pm 2.34	0.40
[La-] (mmol·l ⁻¹)	7.93 \pm 1.92	8.63 \pm 1.49	0.71
50 m RPE	15.14 \pm 1.22	15.86 \pm 1.07	0.24
Stroke Frequency (Hz)	0.81 \pm 0.04	0.81 \pm 0.08	0.79
Stroke Length (m c ⁻¹)	1.89 \pm 0.12	1.87 \pm 0.14	0.74
Stroke Index (m ² c ⁻¹ s ⁻¹)	2.91 \pm 0.34	2.85 \pm 0.31	0.40

Note: WU = Warm-up. [La-] = Blood lactate concentration. RPE = ratings of perceived exertion

Figure 1 shows the percentage differences between the individual times performed without warming up and with previous warm-up. Positive values means better results in the 50 m freestyle after warm-up, whereas negative values means that the swimmers had registered superior 50 m performances without warming up. The results revealed that three of the participants had enhanced performances with warm-up, and the other four without any warm-up procedures.

Figure 1 Percentages of the individual differences between 50m times with no warm-up (No WU) and after regular warm-up (WU) (n=7).



DISCUSSION

The purpose of this study was to evaluate the effect of the warm-up in the 50 m swimming performance, in female national level swimmers. The option for the 50 m freestyle was because it is the shortest event in swimming competition and it is easier to apply and to obtain maximal performances of the swimmers. Main results suggested the 50 m freestyle is not influenced by the previous execution of regular warm-up procedures. There were no statistical differences between the two experimental conditions (with and without warm-up) in the several parameters assessed as the swimming time, [La-], RPE, SF, SL and SI.

Warm-up is used to maximize athlete's performance, increasing muscle and tendon mobility, stimulating blood flow and increase muscle temperature (Smith, 2004). Although the great importance placed in warm-up procedures, it is a fact that their effects or even their ideal structure or type are not well known. Specifically in swimming the limited existing literature is controversy (Fradkin, Zaryn, & Smoliga, 2010). De Vries (1959) and Thompson (1958) suggested improvements in the swimming velocity in short distances (until 91 m) with the previous warm-up. These results were confirmed by the research of Romney and Nethery (1993) that verified significant improvements in the 100 yards performances, after warming up for 15 min. The studies of Mitchell and Huston (1993) and Bobo (1999) reinforced the debate on this matter, as they observed no differences in performance when preceded by different warm-up procedures including no warm-up condition.

In the present research, no differences in the experimental test performed in the two evaluated conditions were observed (Table 1). The 50 m times in freestyle swimming remained equivalent with or without performing the regular warm-up (33.05 ± 2.34 s and 32.71 ± 2.07 s; $p = .24$). In the same way, 25 m partials were similar in the warm-up and in the no warm-up conditions (1st 25 m: 15.01 ± 0.90 s and 15.24 ± 1.06 s; $p = .31$; 2nd 25 m: 17.80 ± 1.33 s and 17.70 ± 1.18 s ; $p = .73$). Similar results were presented by Neiva et al. (2012) comparing the same test conditions in male swimmers. Although these results suggest that regular warm-up is not essential to 50 m performance, Balilionis et al. (2012) verified different results that could lead to opposite conclusions. These authors noticed that regular warm-up led to improvements (~ 0.2 seconds) in 50 yard times when comparing with no warm-up. Complementarily, Neiva et al. (2011) verified that the swimmers exert greater values of maximal and mean propulsive force (11% and 15%, respectively) during 30 seconds of tethered swimming, when previous regular warm up is completed. However, the same research did not show differences in [La-] and in RPE values after the maximal test when preceded by warm-up and with no warm-up common activities. Considering that the trial test performed approaches to 30 s at maximal intensity, the contribution of anaerobic metabolism is essential to fulfill the total energy expenditure (Gastin, 2001). Since [La-] had been commonly used to estimate the contribution of glycolytic metabolism to exercise (di Prampero & Ferretti, 1999), the values seem to highlight the preponderance of the anaerobic

system to meet the energy requirements of the exercise. Mandegue et al. (2005) and Beedle and Mann (2007) proposed that warm-up could be used to maintain the acid-base balance at an appropriated level by stimulating the buffering capacity. This way, it could result in the reduction in the [La-] values. Several studies confirmed these suggestions by observing reductions in blood and muscular lactate concentrations when exercising after priming activities routines (i.e., Gray & Nimmo, 2001; Robergs, Pascoe, Costill, & Fink, 1991). Diverging from these statements, the metabolites assessed in this study kept similar in the two situations. As it can be noticed in Table 1, the non-existence of differences in [La-] values after the 50 m in both conditions corroborated previous results abovementioned, which did not show modifications of the physiological parameters with and without warm-up (De Bruyn-Prevost and Lefebvre, 1980; Neiva et al., 2011; Neiva et al., 2012).

In agreement with previous studies, the level of perceived exertion demonstrated no differences between the two experiments implemented. The RPE values obtained are similar to those presented in previous studies (Neiva et al., 2011; Neiva et al 2012; Ballionis et al., 2012). This scale is used to quantify, monitor and assess an individual's exercise level of exertion (Borg, 1998). Robertson et al. (1986) suggested that an increase in the ratings of perceived exertion could be a consequence of the more usage of anaerobic capacity. The accumulation of ions of hydrogen in the active muscles and blood, follow-on the dissociation of lactic acid can cause a superior perception of effort. As [La-] values did not change in the 50 m performed with warm-up and without warm-up, it would be expected that the RPE values were maintained similar.

Regarding to the stroke cycle kinematics, the registered values showed no differences with and without previous warm-up procedures. SL and SF are independent variables that are related to swimming velocity (Pendergast et al., 2006). The female swimmers of this study did not perform differently the 50 m freestyle in the two test situations, and the swimming velocity was also similar. This could lead to no changes in stroke kinematics as well as in SI. SI is considered as an estimator for overall swimming efficiency (Costill et al., 1985). Thereby, it is assumed that at a given velocity, the swimmer with greater SL has the most efficient swimming. In this case, it can be suggested that warming up did not influence the swimmer efficiency.

Although there was found no statistical differences, it can be noted in Figure 1 that there were swimmers responding differently in each situation. They swim as individual subjects and their performance is also individual, not responding to the warm-up procedures in the same way. In Figure 1 it can be noted that more than 50 % of the female swimmers responded positively to the inexistence of warm-up. However, a great variability was found between swimmers, thus suggesting that coaches should focus their attention on the response of each swimmer, instead of applying the same warm up procedure for all the swimmers.

In conclusion, although the usual changes attributed to warm-up, its efficiency in swimmers performance is still not clear. It was observed that there were no differences in perfor-

mance as well as no differences in the assessed parameters (physiological and biomechanical). These results allow us to suggest that regular warm-up does not seem to be determinant in short distance performances, in female swimmers. Bishop (2003b) pointed some possible explanations for the negative or unaltered results in performance, as the warm-up being of low intensity and not causing the necessary changes in the subjects; being too hard and causing fatigue; and not allowing to recovery sufficiently before exercise. Insufficient and unclear data published, enhances the necessity for further studies to better understand this issue.

PRACTICAL APPLICATIONS

The small sample of subjects does not allow strong statements and certainties, whereby it is needed to amplify the number of swimmers evaluated. Results of the present study indicated no significant influence of using regular warm-up in 50 m swimming, in females. However, as Balilionis et al. (2012) suggested, coaches should work individually with the swimmers, in order to maximize their training or competition performance. It could be noted different responses to the use or no use of warm-up before the maximal trial. This should be taken in consideration when preparing competition, and coaches should focus in each individual needs to achieve optimal performance.

ACKNOWLEDGEMENTS

The authors would like to thank all the swimmers who participated in the study and the Varzim Lazer EEM for allowing using the swimming pool facilities during the experiments. Portuguese government supported this work by a grant of the Science and Technology Foundation (SFRH/BD/74950/2010). This work was also supported by University of Beira Interior and Santander Totta bank (UBI/FCSH/Santander/2010).

REFERENCES

- Atkinson, G., Todd, C., Reilly, T., & Waterhouse, J. (2005). Diurnal variation in cycling performance: influence of warm-up. *Journal of Sports Sciences*, 23(3), 321-9.
- Balilionis, G., Nepocatyč, S., Ellis, C.M., Richardson, M.T., Neggess, Y.H., Bishop, P.A. (2012). Effects of Different Types of Warm-Up on Swimming Performance, Reaction Time, and Dive Distance. *Journal of Strength and Conditioning Research*. doi: 10.1519/JSC.0b013e318248ad40 (in press).

- Beedle, B. B., & Mann, C. L. (2007). A comparison of two warm-ups on joint range of motion. *The Journal of Strength & Conditioning Research*, 21(3), 776-779.
- Bishop, D. (2003a). Warm up I: potential mechanisms and the effects of passive warm up on exercise performance. *Sports Medicine*, 33(6), 439-454.
- Bishop, D. (2003b). Warm up II: performance changes following active warm up and how to structure the warm up. *Sports Medicine*, 33(7), 483-498.
- Bishop, D., Bonetti, D., & Dawson, B. (2001). The effect of three different warm-up intensities on kayak ergometer performance. *Medicine & Science in Sports & Exercise*, 33(6), 1026-1032.
- Bobo, M. (1999). The effect of selected types of warm-up on swimming performance. *International Sports Journal*, 3(2), 37-43.
- Borg, G. (1998). *Borg's perceived exertion and pain scales*. Champaign, IL: Human Kinetics.
- Buchthal, F., Kaiser, E., Knappes, G. G. (1944). Elasticity, viscosity and plasticity in the cross striated muscle fibre. *Acta Physiologica Scandinavica*, 8 (1), 16-37.
- Burnley, M., Doust, J. H., & Jones, A. M. (2002). Effects of prior heavy exercise, prior sprint exercise and passive warming on oxygen uptake kinetics during heavy exercise in humans. *European Journal of Applied Physiology*, 87(4-5), 424-432.
- Costill, D.; Kovaleski, J.; Porter, D.; Fielding, R. & King, D. (1985). Energy expenditure during front crawl swimming: predicting success in middle-distance events. *International Journal of Sports Medicine*, 6, pp. 266-270
- Craig, A. B., Skehan, P. L., Pawelczyk, J. A., & Boomer, W. L. (1985). Velocity, stroke rate, and distance per stroke during elite swimming competition. *Medicine and Science in Sport*, 17(6), 625-634.
- De Bruyn-Prevost, P., Lefebvre, F. (1980). The effects of various warming up intensities and durations during a short maximal anaerobic exercise. *European Journal of Applied Physiology and Occupational Physiology*, 43 (2), 101-7.
- De Vries, H. A. (1959). Effects of various warm-up procedures on 100-yard times of competitive swimmers. *Research Quarterly*, 30, 11-22.
- Di Prampero, P. E., & Ferretti, G. (1999). The energetics of anaerobic muscle metabolism: a reappraisal of older and recent concepts. *Respiration Physiology*, 118(2-3), 103-115.
- Febbraio, M. A., Carey, M. F., Snow, R. J., Stathis, C. G., & Hargreaves, M. (1996). Influence of elevated muscle temperature on metabolism during intense, dynamic exercise. *American Journal of Physiology*, 271(5 Pt 2), R1251-R1255.
- Fradkin, A. J., Zaryn, T. R., & Smoliga, J. M. (2010). Effects of warming-up on physical performance: A systematic review with meta-analysis. *Journal of Strength & Conditioning Research*, 24(1), 140-148.
- Gastin, P. B. (2001). Energy system interaction and relative contribution during maximal exercise. *Sports Medicine*, 31(10), 725-41.
- Gray, S. C., & Nimmo, M. A. (2001). Effects of active, passive or no warm-up on metabolism and performance during short-duration high-intensity exercise. *Journal of Sport Sciences*, 19, 693-700.
- Hedrick, A. (1992). Physiological responses to warm up. *National Strength and Conditioning Journal*, 14(5), 25-27.
- Houmard, J. A., Johns, R. A., Smith, L. L., Wells, J. M., Kobe, R. W., & McGoogan, S. A. (1991). The effect of warm-up on responses to intense exercise. *International Journal of Sports Medicine*, 12(5), 480-483.

- Mandengue, S. H., Seck, D., Bishop, D., Cissé, F., Tsala-Mbala, P., & Ahmaidi, S. (2005). Are athletes able to self-select their optimal warm up? *Journal of Science and Medicine in Sport*, 8(1), 26-34.
- McCutcheon, L. J., Geor, R. J., & Hinchcliff, K. W. (1999). Effects of prior exercise on muscle metabolism during sprint exercise in horses. *Journal of Applied Physiology*, 87(5), 1914-1922.
- Mitchell, J. B., & Huston, J. S. (1993). The effect of high- and low-intensity warm-up on the physiological responses to a standardised swim and tethered swimming performance. *Journal of Sports Sciences*, 11(2), 159-65.
- Neiva, H. P., Morouço, P. G., Pereira, F. M., & Marinho, D. A. (2012). The effect of warm-up in 50 m swimming performance. *Motricidade*, 8(S1), 13-18.
- Neiva, H., Morouço, P., Silva, A. J., Marques, M. C., & Marinho, D. A. (2011). The Effect of Warm up on Tethered Front Crawl Swimming Forces. *Journal of Human Kinetics*, (Special Issue), 113-119
- Pearson, J., Low, D. A., Stöhr, E., Kalsi, K., Ali, L., Barker, H., & González-Alonso, J. (2011). Hemodynamic responses to heat stress in the resting and exercising human leg: insight into the effect of temperature on skeletal muscle blood flow. *American journal of physiology Regulatory integrative and comparative physiology*, 300(3), R663-R673.
- Pendergast, D.R.; Capelli, C.; Craig, A.B.; di Prampero, P.E.; Minetti, A.E.; Mollendorf, J.; Termin, J.I. & Zamparo, P. (2006). Biophysics in swimming, In: *Biomechanics and Medicine in Swimming X*, J.P. Vilas-Boas, F. Alves & A. Marques (Eds.), 185-189, Portuguese Journal of Sport Science, Porto
- Robergs, R. A., Costill, D. L., Fink, W. J., Williams, C., Pascoe, D. D., Chwalbinska-Moneta, J., & Davis, J. A. (1990). Effects of warm-up on blood gases, lactate and acid-base status during sprint swimming. *International Journal of Sports Medicine*, 11(4), 273-278.
- Robergs, R. A., Pascoe D. D., Costill, D. L., & Fink, W. J. (1991). Effect of warm-up on muscle glycogenolysis during intense exercise. *Medicine & Science in Sports & Exercise*, 23, 37-43.
- Robertson, R., Falkel, J., Drash, A., Swank, A., Metz, K., Spungen, S., LeBoeuf, J. (1986). Effects of blood pH on peripheral and central signals of perceived exertion. *Medicine & Science in Sports & Exercise*, 18, 114-122
- Romney, N. C., & Nethery, V. M. (1993). The effects of swimming and dryland warm-ups on 100-yard freestyle performance in collegiate swimmers. *Journal of Swimming Research*, 9, 5-9.
- 94 • Saltin, B., Gagge, A. P., & Stolwijk, J. A. J. (1968). Muscle temperature during submaximal exercise in man. *Journal of Applied Physiology*, 25(6), 679-688.
- Smith, C. A. (2004). The warm-up procedure: To stretch or not to stretch – A brief review. *The Journal of Orthopaedic and Sports Physical Therapy*, 19, 12-17.
- Thompson, H. (1958). Effect of warm-up upon physical performance in selected activities. *Research Quarterly*, 29(2), 231-246.
- Woods, K., Bishop, P., & Jones, E. (2007). Warm-up and stretching in the prevention of muscular injury. *Sports Medicine*, 37(12), 1089-1099
- Wright, V. (1973). Stiffness: a review of its measurement and physiological importance. *Physiotherapy*, 59(4), 107-111.